



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

PROBLEMS AND SOLUTIONS.

SEND ALL COMMUNICATIONS ABOUT PROBLEMS TO B. F. FINKEL, Springfield, Mo.

PROBLEMS FOR SOLUTION.

ALGEBRA.

487. Proposed by WILLIAM HOOVER, Columbus, Ohio.

Show in two ways that 0.5623 is not a root of $(1 - m)e^m = 2e^{-1}$, e being the Napierian base. Find the correct value of m , and test the result in two ways.

488. Proposed by NORMAN ANNING, Chilliwack, B. C.

Show that

$$\begin{vmatrix} a_1 & 1 & a_2 \\ 1 & 1/a_4 & 1 \\ a_2 & 1 & a_3 \end{vmatrix} = 0,$$

where

$$a_k = \frac{\sin(k\theta + \alpha)}{\sin k\theta},$$

and θ and α have any values that do not make a denominator zero.

GEOMETRY.

520. Proposed by ALBERT A. BENNETT, University of Texas.

On a given tangent to a circle determine a point such that, if a secant be drawn joining this point to the extremity of the diameter which is perpendicular to the given tangent, the segment of this secant exterior to the circle will be equal in length to a given segment.

521. Proposed by R. M. MATHEWS, Riverside, California.

A variable circle, with center on the line l and passing through a fixed point P , cuts a fixed circle in A and B . Prove that the common chord AB and the perpendicular to l through P intersect in a fixed point.

CALCULUS.

435. Proposed by B. F. FINKEL, Drury College.

Show that

$$\int_0^\infty e^{-x^2 - (a^2/x^2)} dx = \frac{\sqrt{\pi}}{2e^{a^2}}$$

by a transformation, rather than by the usual method of differentiating under the sign of integration, as, for example, in Byerly's *Integral Calculus*, pages 106-107.

436. Proposed by ARTEMAS MARTIN, LL.D., Washington, D. C.

A circle of radius a is drawn at random on a circular slate of radius r . If another circle of radius a be drawn on the slate, what is the probability that the second circle will intersect the first?

MECHANICS.

352. Proposed by C. N. SCHMALL, New York City.

A glass rod is balanced partly in and partly out of a cylindrical tumbler, with its lower end resting against the vertical wall of the tumbler. If ϕ and ψ are the maximum and minimum angles, respectively, which the rod can make with the vertical plane, and θ is the angle of friction, show that

$$\theta = \frac{1}{2} \tan^{-1} \left(\frac{\sin^2 \phi - \sin^2 \psi}{\sin^2 \phi \cos \phi + \sin^2 \psi \cos \psi} \right).$$